

# PATIENT POSITIONING

## NGN NCLEX MASTER REVIEW · FUNDAMENTALS OF NURSING

Therapeutic positions, clinical indications, contraindications, and high-yield NCLEX patterns. From Fowler's variations to emergency air-embolism positioning — every angle, every rationale, every red flag.

FUNDAMENTALS

SAFETY

OXYGENATION

PROCEDURAL

EMERGENCY

POSTOPERATIVE

CLINICAL JUDGMENT



Critical /  
Emergency



Pathophysiology



Safe / Expected



Complications



NCLEX Trap



Procedure



Pt. Education

## 01 PHYSIOLOGICAL PRINCIPLES OF POSITIONING

How body position alters physiology — the four foundational mechanisms NCLEX tests:

### MECHANISM 01

#### DIAPHRAGM EXCURSION

Upright positions let abdominal contents drop away from the diaphragm → ↑ lung expansion → ↑ tidal volume → ↑ SpO<sub>2</sub>. Critical in ascites, obesity, pregnancy, COPD.

### MECHANISM 02

#### GRAVITY & FLUID SHIFT

Trendelenburg shifts blood centrally; reverse Trendelenburg shifts blood peripherally. Affects BP, ICP, edema, and surgical access to body cavities.

### MECHANISM 03

#### ASPIRATION RISK

Supine + ↓ LOC = pooled secretions → aspiration pneumonia. HOB ↑ 30° minimum protects airway in restraints, post-op, tube feeds, intubated patients.

### MECHANISM 04


#### ALVEOLAR RECRUITMENT

Prone positioning recruits dorsal alveoli collapsed by gravity — improves V/Q matching in ARDS. Frequent rotation prevents pressure injury & atelectasis.

## 02 THE 13 POSITIONS — VISUAL ATLAS

## SEMI-FOWLER'S

HOB 30°–45°




Lying on back; head & trunk elevated 30–45°.

### ✓ INDICATIONS

- Ascites / cirrhosis (↓ diaphragm pressure)
- Restraint use (↓ aspiration risk)
- Post-op thyroidectomy/mastectomy (↓ swelling)
- NG tube feedings (↓ reflux)

## HIGH FOWLER'S

HOB 60°–90°



Nearly upright; maximum lung expansion.

### ✓ INDICATIONS


- Acute dyspnea / SOB
- CHF exacerbation (↓ preload)
- Severe COPD / asthma attack
- Tube feedings (highest aspiration protection)

## ORTHOPNEIC

In bed • over table

## FOWLER'S

HOB 45°–60°




Standard "sitting up in bed" — comfort + oxygenation.

### ✓ INDICATIONS

- General comfort, eating, visiting
- Cardiac/respiratory conditions (↑ lung expansion)
- Post-op recovery (general)
- Support knees with pillow to ↓ shearing

## TRIPOD

Seated • forward lean




Sitting, leaning forward, weight on elbows/knees.

### ✓ INDICATIONS

- COPD exacerbation (often spontaneous)
- Severe asthma attack
- Pneumonia with respiratory distress
- **Sign of distress** if pt assumes spontaneously

## TRENDELENBURG

Whole bed tilted • feet ↑




In bed, leaning forward over bedside table with pillows.

#### ✓ INDICATIONS

- Severe dyspnea (orthopnea)
- Pulmonary edema
- End-stage COPD
- Patients too weak to sit in chair

## REVERSE TRENDLENBURG

Whole bed tilted • head ↑



Pt remains **flat**; entire bed tilted — head higher.

#### ✓ INDICATIONS


- Post femoral cardiac cath (no waist flexion)
- Head/neck/spinal surgery
- ↓ GERD without bending hips

#### ✗ CONTRAINDICATIONS

- Hypotension / hypovolemia
- Peripheral edema, venous stasis

## LEFT LATERAL TRENDLENBURG

EMERGENCY • air embolism



Supine; entire bed tilted — feet higher than head.

#### ✓ INDICATIONS

- Pelvic / lower abdominal surgery (better access)
- Boosting patient up in bed (with lift sheet)

#### ✗ CONTRAINDICATIONS

- ↑ ICP / head injury / stroke
- Ascites, respiratory distress
- GERD, hiatal hernia

## MODIFIED TRENDLENBURG (SHOCK)

Supine • legs elevated 15°–30°



Body flat; **only legs** elevated 15°–30°.


#### ✓ INDICATIONS

- Hypovolemic shock (↑ central venous return)
- Syncope / fainting episode
- Severe hypotension (temporary)

**NOTE:** True Trendelenburg no longer recommended for shock — use modified version.

## SUPINE

Flat • on back




Trendelenburg + **turned to LEFT side** — traps air bubble in right ventricle.

#### ✓ INDICATIONS

- Suspected air embolism (CVC, dialysis line)
- Prevents air from reaching pulmonary artery
- Buys time while team intervenes

**EMERGENCY ONLY · TEMPORARY**



On back, completely flat (no HOB elevation).

#### ✓ INDICATIONS

- Spinal anesthesia recovery
- Post femoral cardiac cath (clot protection)
- Lumbar puncture recovery
- Sleep / rest

#### ✗ CAUTIONS

- ↑ Aspiration risk — use side-lying for unconscious
- ↑ Pressure injury (sacrum, heels)

## PRONE

Flat · on abdomen



On abdomen, head turned to side.

#### ✓ INDICATIONS

- **ARDS** (recruits dorsal alveoli)
- Post lower-limb amputation (prevent hip contracture)
- Post tonsillectomy in children (drainage)

#### ✗ CONTRAINDICATIONS

- Spinal surgery, recent abdominal surgery
- Severe respiratory distress (unmonitored)

## SIMS' (SEMI-PRONE)

Left side · top knee flexed




Semi-prone on LEFT side; right knee flexed forward.

#### ✓ INDICATIONS

- **Enema administration** (eases fluid retention)
- Rectal exam / suppository
- Unconscious patient (drainage of secretions)
- Post-laparoscopy shoulder pain relief

## LITHOTOMY

### Supine • feet in stirrups



Supine; legs flexed at hips & knees, feet in stirrups.

#### ✓ INDICATIONS

- Pelvic / vaginal exam
- Childbirth, cervical procedures
- Cystoscopy (bladder visualization)
- Some perineal surgical procedures

#### ✗ CAUTIONS

- ↑ Risk of nerve injury, DVT — minimize duration
- Raise/lower legs **simultaneously**

## 03 EXPECTED RESPONSE VS CRITICAL FINDINGS

### ✓ EXPECTED — POSITION IS WORKING

- ✓ SpO<sub>2</sub> ↑ ≥ 94% after High Fowler's in dyspneic patient
- ✓ RR returns to 12–20 after orthopneic/tripod for COPD
- ✓ BP stabilizes after modified Trendelenburg in syncope
- ✓ ↓ Reported dyspnea within 2–5 minutes of repositioning
- ✓ Even chest rise, no accessory muscle use post-position change
- ✓ Improved P/F ratio after prone positioning in ARDS (typically 16+ hrs)
- ✓ No bleeding at femoral cath site after supine maintenance
- ✓ Enema fluid retained ≥ 5–10 min in left Sims' position

### ✗ CRITICAL — STOP / REPOSITION / NOTIFY HCP

- ✗ SpO<sub>2</sub> ↓ < 90% after position change → return to prior position
- ✗ ↑ ICP signs (Cushing's triad, ↓ LOC) in Trendelenburg
- ✗ New dyspnea / cyanosis in supine pt → ↑ HOB or side-lying
- ✗ Coughing / gurgling in flat pt with NG feedings → STOP feed, ↑ HOB 30°+
- ✗ Hematoma at cath site if pt flexes hip post-femoral cath
- ✗ Numbness / tingling in legs during prolonged lithotomy
- ✗ Seizure / sudden ↓ LOC → side-lying to protect airway
- ✗ Sudden chest pain + dyspnea in CVC pt → suspect air embolism, L

✓ **Calm, oriented patient** — comfort + ability to communicate needs

lateral Trendelenburg STAT

✗ **Respiratory arrest in prone pt** → emergency supine flip, code algorithm

## 04 ASSESSMENT VALUES & NCLEX SIGNIFICANCE

| PARAMETER                             | EXPECTED VALUE           | CRITICAL                             | NCLEX SIGNIFICANCE                                                                                                    |
|---------------------------------------|--------------------------|--------------------------------------|-----------------------------------------------------------------------------------------------------------------------|
| SpO <sub>2</sub> (positioning effect) | 95–100%                  | < 90%                                | ↑ HOB is the FIRST intervention for SpO <sub>2</sub> < 92% before calling RT or starting O <sub>2</sub> in stable pt. |
| RR                                    | 12–20/min                | < 10 or > 30                         | Tachypnea + tripod posture = severe respiratory distress. Don't waste time — sit pt up immediately.                   |
| HOB angle (tube feeding)              | ≥ 30°                    | < 30°                                | Aspiration prevention standard for ALL enteral feedings, intubated, NG tube patients.                                 |
| ICP                                   | 5–15 mmHg                | > 20 mmHg                            | HOB ↑ 30° in TBI/stroke. NEVER Trendelenburg. Keep neck midline (no rotation).                                        |
| BP (orthostatic)                      | SBP drop < 20 mmHg       | SBP drop > 20 mmHg                   | Position changes (lying → sit → stand) must be SLOW, especially in elderly or post-bedrest.                           |
| P/F ratio (ARDS)                      | > 300                    | < 200                                | Prone for ≥ 16 hrs/day improves P/F ratio in moderate–severe ARDS. Rotate to prevent injury.                          |
| HR (positional)                       | 60–100 bpm               | ↑ > 20 bpm w/ standing               | Reflex tachycardia on standing suggests volume depletion — keep supine, ↑ fluids before mobilizing.                   |
| Femoral cath site                     | Dry, intact, no hematoma | Active bleeding / expanding hematoma | Maintain supine + extended leg × 4–6 hrs. NO hip flexion. Reverse Trendelenburg permitted.                            |

## 05 PRIORITY NURSING ACTIONS — POSITIONING LADDER

01

## ASSESS BEFORE YOU MOVE

Check VS, SpO<sub>2</sub>, LOC, pain, lines/tubes, and contraindications (recent surgery, ↑ ICP, fractures, spinal precautions). NEVER reposition without knowing the order set.

02

## SECURE THE AIRWAY FIRST

If pt is dyspneic → SIT THEM UP (High Fowler's or tripod) before anything else. Airway always wins. Do not delay to take vitals or notify HCP first.

03

## PROTECT FROM ASPIRATION

HOB ≥ 30° for tube feedings, restraints, sedation, post-anesthesia, and any ↓ LOC. For unconscious pt: side-lying (recovery) position.

04

## REPOSITION Q2H MINIMUM

Pressure injury prevention: turn every 2 hours, more often for high-risk (Braden < 18, immobile, incontinent). Document each turn. Use heel offloaders, pillows for bony prominences.

05

## USE PROPER BODY MECHANICS & LIFT EQUIPMENT

Two-person lifts, mechanical lifts (Hoyer), friction-reducing slide sheets. Bend at knees, not back. The most common nursing injury is from improper transfers.

06

## RE-ASSESS AFTER REPOSITIONING

Recheck SpO<sub>2</sub>, RR, BP, pain, and patient comfort within 5 minutes. Position is an INTERVENTION — evaluate its effectiveness and document response.

07

## ADVOCATE & REFUSE UNSAFE ORDERS

If an order conflicts with patient safety (e.g., Trendelenburg in TBI patient), STOP, document, and notify the provider. The nurse is the last line of defense.

06 IF → THEN CLINICAL DECISION RULES

|                                                                  |                                                                                                                              |                                                                    |                                                                                                                     |
|------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------|
| <b>IF</b><br>Patient with CHF develops sudden dyspnea & crackles | <b>THEN</b><br>High Fowler's + dangle legs (orthopneic) → ↓ preload, ↑ lung expansion. Then call HCP for diuretic.           | <b>IF</b><br>Patient is unconscious / sedated / seizing            | <b>THEN</b><br>Side-lying (recovery / lateral) position to protect airway from aspiration. NEVER flat supine.       |
| <b>IF</b><br>Suspected air embolism (CVC removal, dialysis line) | <b>THEN</b><br>LEFT lateral Trendelenburg STAT. Call rapid response. 100% O <sub>2</sub> . Clamp line.                       | <b>IF</b><br>Patient with ↑ ICP / TBI / hemorrhagic stroke         | <b>THEN</b><br>HOB ↑ 30°, neck MIDLINE (no flexion or rotation). NEVER Trendelenburg.                               |
| <b>IF</b><br>Pt s/p femoral cardiac catheterization              | <b>THEN</b><br>Supine, leg straight, NO hip flexion × 4–6 hrs. Reverse Trendelenburg OK to drink fluids.                     | <b>IF</b><br>ARDS pt with refractory hypoxemia despite ventilation | <b>THEN</b><br>Prone position 16+ hrs/day. Rotate to supine briefly. Protect eyes, ETT, lines.                      |
| <b>IF</b><br>Hypotension / syncope / pre-syncope                 | <b>THEN</b><br>Modified Trendelenburg (legs up, body flat). NOT full Trendelenburg.                                          | <b>IF</b><br>Pt admin enema                                        | <b>THEN</b><br>Left Sims' position — anatomic angle of sigmoid colon eases fluid retention.                         |
| <b>IF</b><br>Patient assumes tripod posture spontaneously        | <b>THEN</b><br>RECOGNIZE as severe respiratory distress. Sit pt up, O <sub>2</sub> , call RT, prep for nebulizer/escalation. | <b>IF</b><br>Post-thyroidectomy pt with neck swelling              | <b>THEN</b><br>Semi-Fowler's, neck neutral, trach tray at bedside. Watch for stridor (hematoma airway compression). |
| <b>IF</b><br>Lower-limb amputee at risk for hip contracture      | <b>THEN</b><br>Prone position several times daily to extend hip. Avoid pillow under stump (causes contracture).              | <b>IF</b><br>Tube feeding pt has gastric residual > 250 mL         | <b>THEN</b><br>Hold feeding, maintain HOB ≥ 30°, recheck in 1 hr, notify HCP per protocol.                          |

## 07 ⚠ 8 NCLEX TRAPS — POSITIONING PITFALLS

### "TRENDLENBURG FOR SHOCK"

01

FALSE. Modern guidelines = MODIFIED Trendelenburg (legs up, torso flat). Full Trendelenburg ↑ ICP & ↓ lung expansion.

### "REVERSE TRENDLENBURG = FOWLER'S"

02

NO. Reverse Trendelenburg keeps the body FLAT — the whole bed tilts. Fowler's bends the bed at the waist.

### SUPINE FOR UNCONSCIOUS PT

03

DANGER. Aspiration risk! Use SIDE-LYING (recovery) position for any pt with ↓ LOC, post-seizure, or sedated.

### PILLOW UNDER AMPUTATION STUMP

04

CAUSES CONTRACTURE. Elevate by raising entire foot of bed for first 24 hrs only — then PRONE positioning.

### TRENDLENBURG IN ↑ ICP

05

CONTRAINDICATED — head down ↑ ICP further → herniation risk. Always HOB ↑ 30°, neck midline.

### RIGHT LATERAL FOR AIR EMBOLISM

06

WRONG SIDE. Use LEFT lateral Trendelenburg — traps air in right ventricle, away from pulmonary artery.

### LITHOTOMY: LEGS RAISED SEPARATELY

07

CAUSES hip dislocation / nerve injury. Raise & lower legs SIMULTANEOUSLY with two staff members.

### "ALL COPD PTS NEED O<sub>2</sub> FIRST"

08

POSITION FIRST. Tripod / orthopneic before increasing O<sub>2</sub> in COPD (hypoxic drive). Never push O<sub>2</sub> > 2 L without HCP.

## 08 HIGH-YIELD NCLEX PATTERNS

#### PATTERN 01 • DYSPNEA

"What is the FIRST action for the pt c/o SOB?"

**ANSWER:** Sit the patient upright (High Fowler's). This is BEFORE applying O<sub>2</sub>, calling HCP, or assessing further.

#### PATTERN 02 • POST-OP

"Pt s/p general anesthesia, drowsy. Position?"

**ANSWER:** Side-lying or HOB ≥ 30°. NEVER flat supine — aspiration risk during emergence.

#### PATTERN 03 • NG / TUBE FEEDING

"What HOB angle minimizes aspiration?"

**ANSWER:** HOB ≥ 30° during feeds AND for 30–60 min after. Standard of care for ALL enteral nutrition.

#### PATTERN 04 • ICP

"Position for pt with TBI?"

**ANSWER:** HOB 30°, neck midline (no flexion/rotation). NEVER Trendelenburg.

#### PATTERN 05 • CARDIAC CATH

"Femoral approach — pt position?"

**ANSWER:** Supine, leg extended, HOB ≤ 30° × 4–6 hrs. NO hip flexion.

#### PATTERN 06 • ENEMA

"Position for cleansing enema?"

**ANSWER:** LEFT Sims' position. Anatomic angle of sigmoid eases fluid into colon and improves retention.

#### PATTERN 07 • AIR EMBOLISM

#### PATTERN 08 • ARDS

#### PATTERN 09 • COPD

"CVC pt suddenly dyspneic + chest pain.

Action?"

**ANSWER:** Clamp line, LEFT lateral Trendelenburg, 100% O<sub>2</sub>, call rapid response.

"Pt with severe ARDS, refractory hypoxia.

Position?"

**ANSWER:** Prone for ≥ 16 hrs/day. Improves V/Q matching by recruiting dorsal alveoli.

"Pt with COPD exacerbation, leaning forward.

Significance?"

**ANSWER:** Tripod = sign of severe distress. Recognize and act — escalate care.

#### PATTERN 10 • PEDIATRIC

"Post-tonsillectomy child position?"

**ANSWER:** Side-lying or prone with head turned — promotes oral secretion drainage; ↓ aspiration of blood.

#### PATTERN 11 • MATERNAL

"Pregnant pt c/o dizziness when supine.

Position?"

**ANSWER:** LEFT lateral side-lying — relieves vena cava compression (supine hypotensive syndrome).

#### PATTERN 12 • CRITICAL CARE

"Hypovolemic shock, BP 78/40. Position?"

**ANSWER:** Modified Trendelenburg (legs ↑, body flat). Begin IV fluids immediately.

## 09 NGN BOW-TIE • CLINICAL JUDGMENT MODEL

**SCENARIO:** 62-yr-old female s/p R femoral cardiac catheterization 2 hrs ago. Reports back pain & nausea. Asks for help to get up to use the bathroom. SpO<sub>2</sub> 96% RA, BP 132/82, HR 84.

### ← ACTIONS TO TAKE

Maintain **supine**, leg extended, HOB ≤ 30°

Offer **bedpan or urinal**

Reposition with **log-roll** technique only

Use **Reverse Trendelenburg** if pt wants head up to drink

Assess cath site Q15min × 1hr

### CONDITION MOST LIKELY

#### STANDARD POST FEMORAL CARDIAC CATHETERIZATION RECOVERY

Goal: prevent femoral artery bleeding → strict bed rest with leg straight × 4–6 hrs.

### PARAMETERS TO MONITOR →

**Cath site:** bleeding, hematoma, bruit

**Distal pulses:** dorsalis pedis, post tibial

**Color, temp, sensation** of affected leg

**VS Q15min × 1hr**, then per protocol

**Pain:** sudden back/groin pain = retroperitoneal bleed

# 10 PATIENT & FAMILY EDUCATION CHECKLIST

## ☑ TEACH BEFORE DISCHARGE / POSITION CHANGE

- |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><input type="checkbox"/> <b>Change positions slowly</b> — sit before standing to prevent orthostatic hypotension, especially after bedrest or with antihypertensives.</p> <hr/> <p><input type="checkbox"/> <b>Sleep with head elevated</b> if you have GERD, sleep apnea, CHF, or COPD — extra pillows or wedge.</p> <hr/> <p><input type="checkbox"/> <b>Tripod position is OK for breathing</b> at home with COPD — sit, lean forward, weight on knees during episodes.</p> <hr/> <p><input type="checkbox"/> <b>Don't bend at the waist</b> after femoral cath × 24 hrs — sit up straight, no straining.</p> <hr/> <p><input type="checkbox"/> <b>Use grab bars</b> when changing position in bathroom — biggest fall-risk zone in the home.</p> <hr/> <p><input type="checkbox"/> <b>Family/caregivers:</b> learn safe transfer technique (gait belt, non-skid shoes, lock wheelchair brakes).</p> <hr/> | <p><input type="checkbox"/> <b>HOB 30°+ during &amp; after meals</b> — prevents reflux and aspiration; remain upright 30–60 min after eating.</p> <hr/> <p><input type="checkbox"/> <b>Reposition every 2 hours</b> if confined to bed — pressure injury prevention; check skin daily for redness.</p> <hr/> <p><input type="checkbox"/> <b>Avoid crossing legs</b> after hip surgery (typically 6 wks) — prevents prosthesis dislocation.</p> <hr/> <p><input type="checkbox"/> <b>Pregnant?</b> Sleep on LEFT side — improves blood flow to baby, prevents maternal hypotension.</p> <hr/> <p><input type="checkbox"/> <b>Call before getting up</b> if dizzy, weak, or after surgery / medications — don't risk a fall.</p> <hr/> <p><input type="checkbox"/> <b>Report:</b> sudden chest pain, dyspnea, leg swelling, calf pain, or fainting — call 911.</p> <hr/> |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

## PDF EXPORT INSTRUCTIONS

To save this infographic as a high-resolution PDF: **Ctrl + P** (Windows) or **⌘ + P** (Mac). In the print dialog: select "**Save as PDF**", set **Scale: 65–70%**, enable **Background graphics**, set **Margins: None** or **Minimum**, paper size **Letter** or **A4**. For best results use Chrome or Edge.

© NGN NCLEX Master Review · Patient Positioning · Fundamentals of Nursing — For educational use. Always follow facility protocols and provider orders.