

MUSCULOSKELETAL SYSTEM • NGN NCLEX 2026

Acute Lumbosacral Strain

Low back pain from soft-tissue injury — the #1 cause of activity limitation in adults under 45

⌚ Onset: Sudden

📅 Resolves: 4–6 weeks

🔄 Self-limiting

🚩 Watch for Red Flags



Definition & Overview



WHAT IS IT & WHY IT MATTERS

- ▶ **Definition:** Acute injury to the muscles, ligaments, tendons, or fascia of the lower back (lumbar L1–L5 and sacral region) resulting in pain, spasm, and decreased mobility.
- ▶ **Most common cause** of low back pain (LBP); typically from sudden forceful movement, improper lifting, twisting, falls, or prolonged poor posture.
- ▶ **Self-limiting:** Symptoms usually resolve within **4–6 weeks** with conservative treatment — no surgery, no imaging required unless red flags present.
- ▶ **Risk factors:** Poor body mechanics, obesity, weak core, sedentary lifestyle, smoking (decreases disc nutrition), occupational lifting, prior episodes.

2

Pathophysiology

STEP-BY-STEP MECHANISM



Lumbar (L1-L5) and sacral regions — the most stress-bearing segments of the spine

3

Signs & Symptoms

EARLY VS LATE + RED FLAG FINDINGS

Early / Acute (first 24–48 hr)

- ▶ Sharp, **localized** lower back pain
- ▶ Pain worsens with movement, eases with rest
- ▶ Visible/palpable muscle spasm
- ▶ Decreased range of motion (ROM)
- ▶ Tenderness on palpation
- ▶ Antalgic posture (leaning to one side)

Subacute / Late (after 48 hr)

- ▶ Stiffness on awakening
- ▶ Dull, aching pain
- ▶ Pain may radiate to **buttocks or upper thigh** (NOT below the knee)
- ▶ Fatigue from sleep disruption
- ▶ Limited daily activity tolerance

RED FLAGS — Report Immediately

- ▶ Pain radiating **BELOW the knee** → suspect disc herniation / sciatica
- ▶ Bowel or bladder incontinence → 🚑 CAUDA EQUINA SYNDROME (surgical emergency)
- ▶ Saddle anesthesia (numbness in perineum/inner thighs) → cauda equina
- ▶ Progressive lower extremity weakness or foot drop
- ▶ Fever + back pain → spinal infection / osteomyelitis
- ▶ Night pain unrelieved by rest / weight loss → malignancy
- ▶ History of trauma, IV drug use, or cancer → workup required



Priority Nursing Interventions

ABCS • SAFETY • PRIORITIZATION

- ▶ **ASSESS** pain using PQRST; perform focused neuro exam (sensation, strength, reflexes, gait) — rule out red flags.
- ▶ **POSITION:** Use **William's position** — semi-Fowler's with knees flexed and pillow under knees; OR side-lying with knees flexed and pillow between knees. This unloads the lumbar spine.
- ▶ **ICE first 24–48 hr** (20 min on / 20 min off) to reduce inflammation and spasm.
- ▶ **HEAT after 48 hr** to relax muscles and increase blood flow.
- ▶ **ENCOURAGE early mobilization** — prolonged bed rest is harmful; limit strict bed rest to a maximum of **1–2 days**.
- ▶ **ADMINISTER** prescribed analgesics, NSAIDs, and muscle relaxants on schedule (do not wait for pain to peak).
- ▶ **TEACH** proper body mechanics and lifting technique before discharge.
- ▶ **MONITOR** for worsening neuro symptoms — escalating pain, new numbness, loss of bowel/bladder control = emergency.

5

Pharmacology

FIRST-LINE, SECOND-LINE & ADJUNCT MEDICATIONS

Drug / Class	Mechanism	Key Side Effects	Nursing Considerations
NSAIDs <i>Ibuprofen, Naproxen, Ketorolac</i>	Inhibit COX → ↓ prostaglandins → ↓ inflammation & pain	GI bleed, peptic ulcer, renal toxicity, ↑ BP, fluid retention	Give with food ; monitor BUN/creatinine; ketorolac max 5 days IV/PO
Acetaminophen <i>Tylenol</i>	Central pain reduction (mechanism not fully known)	Hepatotoxicity (esp. with alcohol)	Max 4 g/day (3 g if liver dz); check all combo meds for hidden APAP
Muscle Relaxants <i>Cyclobenzaprine (Flexeril), Methocarbamol, Carisoprodol</i>	CNS depression → ↓ muscle spasm	Drowsiness, dizziness, dry mouth, urinary retention, anticholinergic effects	No driving / alcohol ; fall risk in elderly; short-term use only (1–2 wk)
Opioids <i>Tramadol, Hydrocodone</i> (short-term)	Bind μ-receptors → ↓ pain perception	Constipation, sedation, respiratory depression, dependence	Severe pain only; stool softener; monitor RR; short course
Topical Agents <i>Lidocaine patch, Capsaicin, Diclofenac gel</i>	Local nerve blockade / counterirritant / topical NSAID	Skin irritation, burning sensation (capsaicin)	Wear gloves applying capsaicin; rotate lidocaine patch; off 12 hr/day

6

NCLEX Test Traps

COMMON WRONG CHOICES VS THE CORRECT ANSWER

✗ THE TRAP

"Maintain strict bed rest for 1 week to allow the back to heal."

✓ THE RIGHT ANSWER

Limit bed rest to **1–2 days max**; encourage early gentle mobilization — prolonged rest causes deconditioning and worsens outcomes.

✗ THE TRAP

"Apply a heating pad immediately to relieve the pain."

✓ THE RIGHT ANSWER

ICE first 24–48 hr (inflammation control), then heat. Heat too early increases swelling and pain.

✗ THE TRAP

"Patient reports new urinary incontinence — give the next dose of muscle relaxant."

✓ THE RIGHT ANSWER

🚨 **STOP — notify HCP immediately.** New bowel/bladder dysfunction = cauda equina syndrome, a surgical emergency.

✗ THE TRAP

"Pain radiating to the buttocks always means sciatica — needs an MRI."

✓ THE RIGHT ANSWER

Pain radiating only to **buttocks/upper thigh** can be from strain. **Below-the-knee** radiation suggests sciatica/herniation.

✗ THE TRAP

"Teach the patient to lift heavy objects by bending at the waist with straight legs."

✓ THE RIGHT ANSWER

Bend at the knees, keep back straight, lift with legs, hold the object close to the body. **Never bend AND twist** at the same time.

7

Patient & Family Education

PREVENTION & SELF-MANAGEMENT AT HOME

 **Lift with your legs** — bend knees, keep back straight, hold load close to body.

 **No bend + twist** — move your feet to turn; pivot, don't rotate at the spine.

 **Strengthen core** — abdominal & back exercises stabilize the spine.

 **Maintain healthy weight** — extra weight increases lumbar load.

 **Sleep position** — side-lying with pillow between knees, OR supine with pillow under knees.

 **Posture & ergonomics** — feet flat, lumbar support, screen at eye level; stand every 30 min.

 **Stop smoking** — nicotine reduces disc nutrition and delays healing.

 **Sensible shoes** — low heels; avoid prolonged high-heel wear.

 **Stay active** — walking, swimming, gentle stretching. Avoid bed rest beyond 1–2 days.

 **Call HCP if:** bowel/bladder changes, leg numbness/weakness, fever, pain below knee, no improvement after 4–6 weeks.

8

Memory Boosters

MNEMONICS & QUICK-RECALL PATTERNS

► "BACK PAIN" — Red Flag Mnemonic

- B** Bowel/bladder dysfunction → cauda equina
- A** Anesthesia (saddle) — numbness in perineum
- C** Constitutional symptoms — fever, weight loss, night sweats
- K** Kancer history (cancer red flag)
- P** Pain at night, unrelieved by rest
- A** Age >50 or <20 with new back pain
- I** IV drug use → infection risk
- N** Neuro deficits — weakness, foot drop, progressive numbness

💪 "LIFT SMART" — Safe Body Mechanics

- L** Legs do the work, not the back
- I** In close — hold load near body
- F** Feet shoulder-width apart for stability
- T** Tighten core / abdominals before lifting
- S** Slow and steady — no jerking motions
- M** Move feet to turn — never twist the spine
- A** Avoid lifting above shoulders
- R** Rest between lifts; ask for help
- T** Tools — use carts, dollies, assistive devices

❄️🔥 "ICE then FIRE" — Temperature Therapy Rule

First 48 hours = ICE (cold reduces inflammation & spasm)

After 48 hours = HEAT (warmth relaxes muscle, ↑ circulation)

20 minutes on / 20 minutes off — always with a barrier (towel) between source and skin



NCJMM Clinical Judgment Scenario

SIX-STEP NEXT GENERATION CLINICAL JUDGMENT MEASUREMENT MODEL

Scenario: A 45-year-old construction worker presents to the ED after lifting a heavy box at work 4 hours ago. He reports sudden onset of sharp lower back pain that worsens with movement. Pain radiates to the right buttock but **not below the knee**. Denies bowel/bladder changes, numbness, or weakness. **Vital signs:** BP 142/88, HR 92, RR 18, T 98.6°F, SpO₂ 98% RA. Pain rated **8/10**. Visible muscle spasm in the right lumbar paraspinal area. Neuro exam intact bilaterally. Reports two prior similar episodes that resolved on their own.



RECOGNIZE CUES

Acute onset, clear mechanism of injury (lifting), localized lumbar pain, visible muscle spasm, pain radiating to buttock only (NOT below knee), intact neuro exam, no bowel/bladder changes, no fever, no saddle anesthesia. Elevated BP and HR. Pain 8/10. History of similar episodes.



ANALYZE CUES

Clinical picture is consistent with acute lumbosacral strain — mechanical mechanism, localized pain, muscle spasm, no neuro deficits, no red flags. Elevated vitals likely reflect pain response (sympathetic activation), not infection or hemorrhage. Recurrent episodes suggest occupational risk factor / poor body mechanics.



PRIORITIZE HYPOTHESES

Primary: Acute lumbosacral strain (highest likelihood). **Rule out:** herniated disc (no below-knee radiation), cauda equina syndrome (no bowel/bladder/saddle symptoms), vertebral fracture (no significant trauma), spinal infection (afebrile). **Highest priority concern:** pain control and continuous monitoring for evolving neuro changes.



GENERATE SOLUTIONS

Multimodal pain control (NSAIDs + muscle relaxant + acetaminophen); position in William's position with knees flexed; apply ice to lumbar area; limit bed rest to 1–2 days; teach proper body mechanics and lifting technique; arrange physical therapy follow-up; provide red-flag warning signs for return to ED.



TAKE ACTION

Administer ordered ibuprofen 600 mg PO and cyclobenzaprine 10 mg PO. Position patient supine with pillow under knees (William's position). Apply ice pack 20 min with towel barrier. Reassess pain in 60 minutes. Initiate body mechanics teaching. Provide written discharge instructions with red flags.



EVALUATE OUTCOMES

At 60 min: pain decreased from 8/10 to 4/10, muscle spasm reduced, BP 128/80, HR 78, able to ambulate cautiously without assistance, neuro exam remains intact, no new red flags. **Interventions effective.**

Patient verbalizes understanding of body mechanics and red-flag symptoms. Discharge home with follow-up in 1 week.

NGN NCLEX 2026 Visual Study Library • Acute Lumbosacral Strain • Musculoskeletal System

Content drawn from standard NGN NCLEX 2026 references (Saunders, ATI, Lippincott) — topic not present in uploaded notes