

• NGN NCLEX 2026 • INFECTIOUS DISEASE / RESPIRATORY

Tuberculosis (*TB*)

A high-yield clinical judgment reference for the bacterial infection you must never miss — airborne precautions, the RIPE regimen, and the reasoning behind every nursing priority.

🟢 **Mycobacterium tuberculosis**

🌬️ **Airborne transmission**

🔥 **Primarily pulmonary**

🕒 **6–12 month therapy**

🚑 **Reportable disease**

1 Definition & Overview

FOUNDATION

WHAT & WHY


An airborne bacterial infection of the lungs

- Caused by the **Mycobacterium tuberculosis** — a slow-growing, waxy-walled acid-fast bacillus
- Transmitted **inhaled** (coughing, sneezing, singing, talking) — **droplet nuclei** *not* by surfaces, food, or blood.
- Can exist as **latent TB infection (LTBI)** — walled off, asymptomatic, not contagious — or **active TB disease**, which IS contagious and must be treated.

1 in 4

~10%

OF LATENT CASES REACTIVATE




PEOPLE WORLDWIDE CARRY LATENT TB

Upper-lobe cavitation & granulomas are the classic pulmonary TB findings on imaging.

2 Pathophysiology

MECHANISM



Latent TB Infection (LTBI)

- PPD/IGRA **POSITIVE**
- Chest x-ray **normal**
- Sputum culture **negative**
- **Not** contagious, no symptoms
- Treat with INH alone × 6–9 months

Active TB Disease

- PPD/IGRA positive + symptoms
- Chest x-ray: upper-lobe infiltrates, cavitation
- Sputum AFB **POSITIVE**
- **IS** contagious — airborne precautions
- RIPE multi-drug therapy × 6–12 months

3 Signs & Symptoms

CLINICAL PRESENTATION

Early / Mild

EASY TO MISS

Late / Severe

RED FLAGS



Low-grade afternoon fever (classic TB timing)



Drenching night sweats — soaks pajamas / sheets



Fatigue, malaise, lethargy



Anorexia & unintentional weight loss



Persistent cough > **3 weeks**



Hemoptysis — blood-tinged / frankly bloody sputum



Productive cough with **mucopurulent sputum**



Dyspnea on exertion



Pleuritic chest pain



Severe wasting — historic name "*consumption*"



PRIORITY CUE: *A cough lasting longer than 3 weeks — especially with night sweats and weight loss — warrants TB screening. Assume airborne isolation before confirmation if suspicion is high.*

4

Priority Nursing Interventions

ABC & SAFETY

A

Airway & Infection Control

- **Airborne precautions** — priority intervention before anything else
- Obtain **3 early-morning sputum** specimens for AFB smear & culture
- Encourage cough & deep breathe to mobilize secretions

B

Breathing & Oxygenation

- Auscultate lungs every shift — crackles in upper lobes
- Monitor SpO₂, respiratory rate, and work of breathing
- Semi-Fowler's position to optimize lung expansion

C

Circulation & Systemic

- Monitor **hemoptysis** — amount, color, for frequency
- Assess nutrition & weight — offer high-protein, high-calorie meals
- Track **LFTs** (RIPE drugs are hepatotoxic) & vision (ethambutol)



Airborne Precautions Checklist

- ✓ **Private, negative-pressure room** (AIIR) with ≥ 6 –12 air exchanges/hr
- ✓ **Nurse wears a fit-tested N95** (or PAPR) respirator before entering
- ✓ **Teach the client to cover cough**, use tissues, perform hand hygiene
- ✓ Keep the **door closed** at all times
- ✓ **Client wears a surgical mask** whenever leaving the room (not N95)
- ✓ Report confirmed cases to the **public health department**

5 Pharmacology — The RIPE Regimen

FIRST-LINE DRUGS

*R*ifampin *I*soniazid *P*yrazinamide *E*thambutol

2 months of all 4 → then 4 months of INH + Rifampin

DRUG	MECHANISM	KEY SIDE EFFECTS	NURSING CONSIDERATIONS
Rifampin R · BACTERICIDAL	Inhibits bacterial RNA synthesis	Hepatotoxicity ; orange/red discoloration of body fluids (urine, saliva, tears, sweat); GI upset; ↓ effectiveness of oral contraceptives , warfarin , digoxin	Reassure that orange fluids are expected — not a reason to stop. Teach backup contraception. Monitor LFTs; avoid alcohol.
Isoniazid (INH) I · BACTERICIDAL	Inhibits mycolic acid cell wall synthesis	Peripheral neuropathy (numbness/tingling); hepatotoxicity ; can cause B6 deficiency	Give vitamin B6 (pyridoxine) daily to prevent neuropathy. Take on empty stomach. Avoid tyramine-rich foods & alcohol. Monitor LFTs.
Pyrazinamide P · BACTERICIDAL	Disrupts bacterial membrane transport	Hepatotoxicity ; hyperuricemia → gout flare ; photosensitivity; arthralgia	Monitor LFTs and uric acid. Encourage fluids. Teach sunscreen & protective clothing. Contraindicated in pregnancy.
Ethambutol E · BACTERIOSTATIC	Inhibits arabinosyl transferase (cell wall)	Optic neuritis — blurred vision, red–green color blindness , decreased visual acuity	Baseline + monthly vision screening (Snellen & color). Teach the client to report any vision change immediately — usually reversible if caught early.

⚠️ **Shared hepatotoxicity warning:** R, I, and P are all hepatotoxic. Baseline and routine **LFTs** are mandatory. Teach the client to report *yellowing of skin/eyes, dark urine, RUQ pain, nausea, or anorexia*. **Absolutely no alcohol** during therapy.

6 NCLEX Test Traps

COMMON MISSES

1

"Positive PPD = active TB" — FALSE

A positive Mantoux only proves **exposure / infection**. Active disease requires a **positive AFB sputum culture** + symptoms + chest x-ray findings. Don't pick "start airborne precautions on a positive PPD alone" without other cues.

2

N95 for the nurse — surgical mask for the client

~~"Give the client an N95"~~ → **"Give the client a surgical mask when leaving the room."** An N95 must be fit-tested; the client can't achieve a proper seal and doesn't need to filter their own exhaled droplets.

3

Orange urine on rifampin = expected, not toxicity

If a client reports orange/red tears, urine, or sweat, **reassure them**. This is a harmless staining effect. Warn them it can permanently stain soft contact lenses. Stopping the drug is the wrong answer.

4

Induration — not erythema — is what you measure

Read the Mantoux at **48–72 hours**. Measure the **raised, hardened area (induration)** in mm — *not* redness. Cut-offs: ≥ 5 mm (HIV/immunocompromised), ≥ 10 mm (high-risk), ≥ 15 mm (low-risk/healthy).

5

Non-infectious criteria — all three must be met

A client is considered non-contagious only when: **(1)** consistent drug therapy ≥ 2 –3 weeks, **(2)** clinical improvement, and **(3)** **three consecutive negative AFB sputum smears**. Missing any of these = still on airborne precautions.

6

Neuropathy on INH isn't a "stop the drug" answer

It's a **"give pyridoxine (B6)"** answer. B6 is co-prescribed prophylactically with INH. The priority is prevention, not discontinuation.

7

Patient Education

TEACH-BACK PRIORITY



Medication Adherence

- ✓ Take **ALL** medications for the **FULL** 6–12 months — stopping early causes resistance (MDR-TB)
- ✓ Expect **Directly Observed Therapy (DOT)** from public health
- ✓ Take INH & Rifampin on an empty stomach when possible
- ✓ Never double a dose if one is missed — follow provider guidance



Home & Family Safety

- ✓ Open windows & maintain airflow — sunlight and ventilation kill TB bacilli
- ✓ Cover mouth/nose with tissue when coughing or sneezing; dispose carefully
- ✓ **All household contacts must be tested** (PPD or IGRA)
- ✓ Sleep in a separate room until cleared as non-infectious



Report Immediately

- ✓ **Vision changes** (ethambutol — optic neuritis)
- ✓ **Yellow skin/eyes**, dark urine, RUQ pain (hepatotoxicity)
- ✓ **Numbness/tingling** of extremities (INH neuropathy)
- ✓ Worsening cough, hemoptysis, or fever despite treatment



Lifestyle & Follow-Up

- ✓ **No alcohol** — dramatically increases liver toxicity risk
- ✓ High-protein, high-calorie diet — combat weight loss
- ✓ Use a backup contraceptive method (rifampin ↓ OCP effect)
- ✓ Keep all appointments: LFTs, sputum cultures, vision checks

8

Memory Boosters

MNEMONICS

FIRST-LINE DRUGS

RIPE

The 4-drug combo you start every active TB case on.

R **Rifampin** — red-orange body fluids; hepatotoxic

I **Isoniazid (INH)** — needs B6; neuropathy; hepatotoxic

P **Pyrazinamide** — hyperuricemia (gout); hepatotoxic

CLASSIC TB TRIAD

The 3 W's

Three signs that should always make you think TB.

W **Weight loss** — unintentional, progressive

W **Wet pillow** — drenching night sweats

E **Ethambutol** — eye issues (optic neuritis)

W **Worsening cough** — persistent > 3 weeks, eventually with hemoptysis

DRUG SIDE-EFFECT RECALL

Sound-alike Cues

Quick associations that stick before exam day.



"Red-Orange Pee" → Rifampin



"Eye-thambutol" → Ethambutol = eye toxicity



"INH Is Not Happy without B6" → give pyridoxine



"Pyrazinamide + uric acid" → think gout

ISOLATION SHORTCUT

TB Room Rules

Six-word recipe for the NCLEX isolation question.

N

Negative—pressure private room

9

N95 fit-tested respirator — for the nurse

5

5 door rule — keep it closed

S

Surgical mask — for the client when leaving

NCJMM Clinical Judgment — *Apply It*

NGN NCLEX 2026

Scenario: A 42-year-old client presents to the clinic with a 4-week productive cough, 15-lb unintended weight loss, and soaked sheets at night. Temp 100.4°F; crackles auscultated in the right upper lobe. PPD induration is 16 mm at 72 hours.

STEP 1 · RECOGNIZE

Recognize Cues

Persistent cough > 3 weeks, weight loss, night sweats, low-grade fever, RUL crackles, positive PPD (≥ 15 mm).

STEP 2 · ANALYZE

Analyze Cues

Cluster matches active pulmonary TB. Positive PPD confirms exposure, not disease — must confirm with chest x-ray and 3 AFB sputum samples.

STEP 3 · PRIORITIZE

Prioritize & Act

PRIORITY: Place client in a **negative-pressure room immediately**; don an N95; obtain sputum $\times 3$; initiate RIPE therapy; notify public health.

STEP 4 · EVALUATE

Evaluate Outcomes

After 2–3 weeks of therapy, expect 3 negative AFB smears, resolving cough, weight stabilization, and normal LFTs. Only then remove airborne isolation.